



Biogeography

Biodiversity, distribution, and conservation status of Pinaceae in Puebla, Mexico

Biodiversidad, distribución y estado de conservación de la familia Pinaceae en Puebla, México

Diana B. Ramos-Dorantes, José Luis Villaseñor, Enrique Ortiz, David S. Gernandt*

Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado Postal 70-367, 04510 Mexico City, Mexico

Received 15 April 2016; accepted 13 November 2016

Available online 23 February 2017

Abstract

Pinaceae is the most species rich conifer family in Mexico. Here we describe the diversity and geographic distribution of the family in the state of Puebla based on revision of herbarium specimens and field exploration. 572 georeferenced records for 15 species and 1 variety were mapped. These were included in potential distribution models to predict the most species-rich regions in the state and quantify the known species distributions in $10' \times 10'$ (latitudinal \times longitudinal) cells. The conservation status of each taxon was evaluated by calculating its area of occupation and conservation category within the state. A species–accumulation curve agreed with the number of taxa employed, suggesting that the family is relatively well sampled. The resulting potential distribution models indicated that the regions of highest predicted diversity in the state correspond well to the actual diversity estimates, although in some sites more species were predicted than were recorded. The Sierra Madre Oriental and the Trans-Mexican Volcanic Belt were the biogeographic regions with the greatest known and potential richness. Finally, the regional evaluation of conservation status provides relevant information for local conservationists and authorities, given that 12 of the 16 taxa were determined to be at risk.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: *Abies*; Conservation status; *Pinus*; Potential distribution; *Pseudotsuga*; Richness

Resumen

Pinaceae es la familia de coníferas de mayor riqueza en México. Aquí se describe la riqueza de las especies y la distribución geográfica de la familia en el estado de Puebla basada en la revisión de ejemplares de herbario y trabajo de campo. Se empleó un total de 572 registros de 16 taxones georreferenciados equivalentes a 15 especies y una variedad de Pinaceae para Puebla. Se utilizaron modelos de distribución potencial para predecir las regiones de mayor riqueza en el estado y conocer las estimaciones actuales dentro de celdas de $10' \times 10'$ (longitud \times latitud). Para saber qué tan adecuado ha sido el muestreo de la familia, se empleó una curva de acumulación de especies. El estado de conservación se obtuvo reportando el área de ocupación de cada especie y su categoría de amenaza dentro del estado. El resultado obtenido con la curva de acumulación permite suponer que en Puebla ha habido una exploración botánica satisfactoria de la familia. Los modelos de distribución potencial obtenidos exponen de manera general un patrón similar al de la riqueza conocida, excepto que en algunos sitios se predice un mayor número de especies de las que actualmente se tiene registrado. En ambos se aprecia que las regiones biogeográficas con la mayor riqueza de especies son el Eje Volcánico Transversal y la Sierra Madre Oriental. Finalmente, la evaluación regional del estado de conservación ofrece información relevante que pudiera ser tomada en cuenta por conservacionistas locales y autoridades, puesto que dentro del territorio 12 taxones tuvieron una categoría de riesgo.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Palabras clave: *Abies*; Estado de conservación; *Pinus*; Distribución potencial; *Pseudotsuga*; Riqueza

* Corresponding author.

E-mail address: dgernandt@ib.unam.mx (D.S. Gernandt).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

Introduction

Conifers are the largest gymnosperm lineage, making up approximately 64% of the group's species richness (Eckenwalder, 2009). Pinaceae is the largest conifer family, comprising approximately 232 species classified in 11 genera (Farjon & Filer, 2013). The family is distributed exclusively in the Northern Hemisphere except for 1 equatorial species of *Pinus* in Sumatra. Most species occur in montane habitats, particularly in conifer forests in temperate to cold regions; less commonly they occur in semi-arid regions and in tropical and subtropical latitudes (Farjon & Filer, 2013).

In Mexico, Pinaceae is represented by 4 genera (*Abies*, *Picea*, *Pinus*, and *Pseudotsuga*) and 61 species, 30 of which are endemic (Gernandt & Pérez-de la Rosa, 2014). Pinaceae genera are the defining components of *Pinus* forest, *Pinus* shrubland, *Abies* forest, and *Pseudotsuga* and *Picea* forest; they also occur in other vegetation types such as montane cloud forest, mixed forest (most notably, *Pinus-Quercus* forest), and xeric shrublands (Rzedowski, 1978). *Pinus* is the most species rich genus in the family, and Mexico has a greater richness and diversity of pines than any other country in the world (Farjon & Styles, 1997).

The state of Puebla is located in central Mexico between 20°50' N and 17°52' N, 96°43' W and 99°04' W. It has an area of 34,290 km², which represents 1.7% of the total area of the country (Inegi, 2004). In the state 7 biogeographical regions converge, of which, the 4 most extensive are the Sierra Madre Oriental, the Northern Gulf Coastal Plain, the Trans-Mexican Volcanic Belt, and the Balsas River Basin (Inegi, 2004). The state possesses a great diversity of climates and vegetation types, due fundamentally to striking altitudinal changes and its relief. Temperate climates predominate, with forests having the greatest extension, and conifer forests being exceptionally widespread (Inegi, 2000). For Puebla, 15 species of Pinaceae are reported, corresponding to 24.6% species diversity of Pinaceae for Mexico (Ramos-Dorantes, 2015; Table 1). These belong to 3 genera: *Abies* (2 species), *Pinus* (the most diverse with 12 species and 4 varieties), and *Pseudotsuga* (1 species). Their distributions are known from herbarium specimens and databases. However, it is uncertain in quantitative terms how adequately the state has been explored. With the help of computational tools and geographic information systems (GIS) it has been possible to evaluate different aspects of conifers, for example to estimate the distribution of species individually (García-Aranda, Cantú-Ayala, Estrada-Castillón, Pando-Moreno, & Moreno-Talamantes, 2012; Leal-Nares et al., 2012) or collectively (Evans & Cushman, 2009). They have also served to evaluate the distribution of species in the past (Gugger, González-Rodríguez, Rodríguez-Correa, Sugita, & Cavender-Bares, 2011) and under climate change scenarios (Gómez-Mendoza & Arriaga, 2007), to estimate niche divergence between species (Aguirre-Gutiérrez, Serna-Chávez, Villalobos-Arámbula, Pérez-de la Rosa, & Raes, 2015; López-Reyes, Pérez-de la Rosa, Ortiz, & Gernandt, 2015), to assess conservation strategies either in the present or in the future (Aguirre-Gutiérrez & Duivenvoorden, 2010; Ogawa-Onishi, Berry, & Tanaka, 2010), and even to model the

Table 1

Pinaceae taxa recorded for the state of Puebla.

<i>Abies</i>	<i>Abies hickelii</i> Flous & Gaussen, <i>Abies religiosa</i> (Kunth) Schldtl. & Cham.
<i>Pinus</i>	<i>Pinus ayacahuite</i> Ehrenb. ex Schldtl. <i>Pinus ayacahuite</i> var. <i>veitchii</i> (Roezl) Shaw* <i>Pinus chiapensis</i> (Martínez) Andersen <i>Pinus cembroides</i> Zucc. <i>Pinus cembroides</i> subsp. <i>orizabensis</i> (D.K. Bailey)* <i>Pinus douglasiana</i> Martínez <i>Pinus greggii</i> Engelm. ex. Parl. <i>Pinus hartwegii</i> Lindl. <i>Pinus leiophylla</i> Schiede ex. Schldtl. & Cham. <i>Pinus montezumae</i> Lamb. <i>Pinus oocarpa</i> Schiede ex Schldtl. <i>Pinus patula</i> Schiede ex Schldtl. & Cham. <i>Pinus pseudostrabus</i> Lindl. <i>Pinus pseudostrabus</i> fo. <i>protuberans</i> Martínez* <i>Pinus pseudostrabus</i> var. <i>apulcensis</i> (Lindl.) Shaw <i>Pinus teocote</i> Schiede ex Schldtl. & Cham.
<i>Pseudotsuga</i>	<i>Pseudotsuga menziesii</i> var. <i>glauca</i> (Beissn.) Franco

* Intraspecific taxa not analyzed in this work.

distribution of their parasites (Evangelista, Kumar, Stohlgren, & Young, 2011). Similar strategies have been applied in other groups of plants (Cruz-Cárdenas, Villaseñor, López-Mata, & Ortiz, 2013; Cruz-Cárdenas, López-Mata, Ortiz-Solorio, et al., 2014; Cruz-Cárdenas, López-Mata, Villaseñor, & Ortiz, 2014; Delgadillo, Villaseñor, Cárdenas, & Ortiz, 2014; Villaseñor, Ortiz, Cadena, & Estrada, 2013). The objective of the present study is to evaluate the distribution, species richness, and conservation status of Pinaceae in the state of Puebla by analyzing data from natural history collections, through field exploration, and by generating potential distribution models.

Materials and methods

We used 572 georeferenced specimen records of 15 species reported by Ramos-Dorantes (2015) as occurring naturally in Puebla (Table 1). *P. pseudostrabus* was separated into 2 varieties (vars. *pseudostrabus* and *apulcensis*), but the same distinction was not made for 3 other infraspecific taxa recorded for the state. Records were obtained from field collections and from 6 herbaria (MEXU, INIF, ENCB, CHAP, CHAPA, and HUAP). We also consulted online collections (The New York Botanical Garden, 2014; The United States National Herbarium, 2014) and databases (SEInet, 2014), and the Global Biodiversity Information Facility (GBIF, 2014).

Most studies of distribution patterns have focused on measuring the number of species in equivalent areas within the study area. For that reason, they have used a constant cell size and counted the number of species whose distribution intersects a given cell (Arita-Watanabe, 2001). This strategy initially identifies cells with high species richness. To estimate the appropriate cell size for the present study, information from the records was used to calculate the area of occupation (AOO) of each species. To define this area, we followed the recommendation by the International Union for the Conservation of Nature (IUCN,

Table 2
Taxon distribution data.

Species	Number of records training/test	Cells with records	AUC	Threshold description	Training omission rate	Test omission rate	AOO (km ²)	AOO category (IUCN, 2001)	Area of potential distribution (km ²)
<i>Abies hickelii</i>	4/1	2	0.999	*	0.000	1.000	108.02	Endangered (EN)	204.908
<i>A. religiosa</i>	30/10	19	0.970	◦	0.100	0.100	4,340.71	Possible Near Threatened (NT)	2,610.782
<i>Pinus ayacahuite</i>	14/4	11	0.938	~	0.143	0.250	2,896.90	Possible Near Threatened (NT)	6,553.203
<i>P. cembroides</i>	24/8	14	0.986	^	0.000	0.125	2,954.43	Possible Near threatened (NT)	1,830.553
<i>P. chiapensis</i>	16/5	10	0.993	*	0.000	0.800	596.08	Vulnerable (VU)	1,051.474
<i>P. douglasiana</i>	4/1	2	0.999	^	0.000	0.000	18.31	Endangered (EN)	392.54
<i>P. greggii</i>	4/1	2	0.954	*	0.000	1.000	3.22	Critically Endangered (CR)	1,207.562
<i>P. hartwegii</i>	22/7	15	0.974	*	0.045	0.429	4,778.75	Possible Near threatened (NT)	2,897.903
<i>P. leiophylla</i>	33/11	19	0.969	*	0.000	0.182	6,262.64	Possible Near Threatened (NT)	5,902.206
<i>P. montezumae</i>	30/10	21	0.977	*	0.000	0.300	6,118.50	Possible Near Threatened (NT)	3,488.706
<i>P. oocarpa</i>	4/1	1	0.995	*	0.000	1.000	0.11	Critically Endangered (CR)	503.447
<i>P. patula</i>	33/11	22	0.963	*	0.030	0.455	10,327.20	Least Concern (LC)	4,131.744
<i>P. pseudostrobus</i>	36/12	30	0.936	^	0.056	0.167	14,760.70	Least Concern (LC)	6,919.601
<i>P. pseudostrobus</i> var. <i>apulcensis</i>	35/11	30	0.943	*	0.086	0.364	11,488.30	Least Concern (LC)	9,897
<i>P. teocote</i>	52/17	35	0.938	"	0.115	0.235	14,742.3	Least Concern (LC)	8,971.177
<i>Pseudotsuga menziesii</i> var. <i>glauca</i>	12/3	8	0.995	*	0.000	0.000	738.73	Vulnerable (VU)	751.853

* Maximum training sensitivity plus specificity.

◦ 10 percentile training presence.

~ Equal training sensitivity and specificity.

^ Maximum test sensitivity plus specificity.

" Equal test sensitivity and specificity.

2001) to set the cell size to 10% of the distance separating the 2 most distant collection points (Suárez-Mota & Villaseñor, 2011). This was done with the “Conservation Assessment Tools” extension for ArcView (Moat, 2007). The optimal width of the cell was the mean of the values of the cell of each species, and the result was a cell of 14.8 km per side. These were transformed to arcminutes, giving a measurement of 10 min. This measurement was used to form a quadrant over the state of Puebla. The geo-referenced Pinaceae collection records were placed over this quadrant to obtain the number of species in each geographic unit (cell) and a map of known richness was generated with this information.

We used the geographic coordinates of the collection records to construct a species–accumulation curve (Gotelli & Colwell, 2001). The sampling units were 70 cells of 10 min per side. The asymptote of the accumulation curve theoretically predicts the number of species that should be found in the study area (Jiménez-Valverde & Hortal, 2003). The number of cells in which a taxon appeared was used as a measurement of the collection effort. Its order was randomized 100 times using

EstimateS version 9.1.0 (Colwell, 2009), with the aim of constructing a smoothed curve. The asymptote of the curve was estimated by adjusting the Clench equation to the accumulation curve (Colwell & Coddington, 1994; Soberón & Llorente, 1993) using the Simplex and Quasi-Newton method in STATISTICA (StatSoft, 2011).

We used 58 environmental variables for Puebla with a resolution of 30 arcseconds (López-Mata, Villaseñor, Cruz-Cárdenas, Ortiz, & Ortiz-Solorio, 2012). At this resolution, we performed a principal component analysis (PCA) to select variables for use in modeling (Cruz-Cárdenas, López-Mata, Villaseñor, et al., 2014; Janžekovič & Novak, 2012). PCA is one of the most commonly used methods for removing correlation from a set of variables and reducing collinearity (Dormann et al., 2013). We chose the components that accumulated 85% of the variance, and for each component the 5 variables with the greatest absolute load value. The analysis was performed with R (R Development Core Team, 2015).

The potential distribution map was generated with Maxent v. 3.3.3k (Phillips, Anderson, & Schapire, 2006) using the

default settings. For training the model 75% of the records were used and the remaining 25% for testing. The full state's surface was considered as the mobility area (M of the BAM schedule of Soberón & Peterson, 2005) for all the species analyzed. The contribution of the variables was estimated using the jackknife. To implement the jackknife, variables were excluded 1 at a time and the model was re-run to compare its performance in describing the species distribution in the absence of the variable (Baldwin, 2009; Phillips et al., 2006). The minimum number of records used to perform the modeling was 5 per species; for those with fewer records, models were generated using georeferenced records from the adjacent states of Hidalgo and Veracruz. Each distribution model was transformed to a Boolean (presence–absence) layer (Peterson et al., 2011) using ArcMap 10 with a different threshold for each species with the aim of minimizing the error or omission rate (Table 2). The potential distribution map obtained for each species indicated the areas in the state where the model predicted its presence (Graham & Hijmans, 2006). Subsequently, the Boolean models were overlaid to generate a potential distribution map for Pinaceae in Puebla.

To estimate the conservation status of each species we followed the categories and criteria of the IUCN Red List (IUCN, 2001). We again used the “Conservation Assessment Tools” extension for Arcview (Moat, 2007) to calculate the area of occupancy (AOO) of each taxon and its risk category within the state of Puebla.

Results

The distribution of Pinaceae is heterogeneous in the state of Puebla. The majority of the cells with records had more than 3 taxa. The cells with the greatest number of species (7 or 8) had 12, and 30 cells had only 1 or 2 (Fig. 1). The most widely distributed species in the state was *P. teocote* (35 cells), followed by *P. pseudostrobus* and *P. pseudostrobus* var. *apulcensis*, both occurring in 30 cells. *Abies hickelii*, *P. douglasiana*, *P. greggii*, and *P. oocarpa* are the rarest species in the state, occurring in only 1 or 2 cells (Table 2).

The species–accumulation curve had an asymptote of 16 (Fig. 2). This estimate coincides with the number of Pinaceae taxa included in the present study.

The PCA of the 58 variables recovered 7 components that accumulated 85% of the variance. The 5 variables with the greatest absolute load values were identified from each of these components, resulting in the selection of 28 variables (Table 3). These variables were used for modeling.

Based on the jackknife test, the variables with the greatest contribution to modeling the distribution in the majority of species were Bio1 (mean = 13.39, standard deviation = 2.95; 10 species), CE (mean = 0.16, standard deviation = 0.10; 8 species), and MO (mean = 2.48, standard deviation = 0.86; 7 species). The remaining variables differed among the species. In some cases, only 1 variable contributed (Table 4). Moreover, Bio9 and Bio18 had almost a null contribution to the modeling.




Figure 1. Known richness of Pinaceae in Puebla ($N = 16$ taxa). The tones indicate the number of species registered in each cell (10 arcminutes per side).




Figure 2. Accumulation curve. The circles represent the sampling units (cells of 10 arcminutes per side). The asymptote reaches 16 species. The parameters of the curve are indicated in the equation in the upper part of the figure.

The species with the smallest potential distribution was *A. hickelii*, whereas that with the largest was *P. teocote* (Table 1; Fig. 3). The potential distribution models of the 16 taxa allowed us to generate a map (Fig. 5). The maximum potential number of taxa predicted for a cell was 10.




Figure 3. Species of Pinaceae with smallest (*Abies hickelii*; left) and largest (*Pinus teocote*; right) potential distributions in Puebla.

The categories of conservation according to the area of occupation ranged from Least Concern to Seriously Threatened (Table 2).

Discussion

The cells with the greatest concentration of known species of Pinaceae in the state of Puebla are concentrated in the principal elevations of the Trans-Mexican Volcanic Belt, specifically Popocatepetl, Iztaccíhuatl, Malinche, and Pico de Orizaba (Fig. 1). This is where temperate climatic conditions favor the distribution of the family (Inegi, 2004). Cells in the Sierra Norte de Puebla also stand out. Generally, species richness of Pinaceae is concentrated along the mountain chains of the Trans-Mexican Volcanic Belt and the Sierra Madre Oriental. These 2 large biogeographic regions harbor the greatest number of gymnosperms, particularly conifers (Contreras-Medina, Castañeda-Aguado, & González-Zamora, 2007; Contreras-Medina & Luna-Vega, 2007). Pinaceae is the best represented in both regions, as the family accounts for more than half of the gymnosperm species distributed there. This agrees with the principal types of vegetation in the state. The greatest percentage of species is found in pine–oak forest, pine forest, and cloud forest. The latter forest type is better represented in the Sierra Madre Oriental (Contreras-Medina & Luna-Vega, 2007; Contreras-Medina et al., 2007). These forests are considered important areas of diversity for gymnosperms, particularly the genus *Pinus* (Contreras-Medina & Luna-Vega, 2007; Contreras-Medina et al., 2007; Farjon & Styles, 1997; Sánchez, 2008).

The cells with the lowest species richness occur adjacent to the coastal region of the Gulf of Mexico, and in the central and southern parts of the state belonging to the Balsas Basin. The Gulf of Mexico is dominated by perennial rainforests or lowland deciduous rainforests, and the Balsas Basin by shrublands (Inegi, 2000). Some cells lacked recorded species, even when nearby cells did have records. This could be explained by the lack of sampling in these areas, by the cells lacking favorable environments for the growth of Pinaceae, or by anthropogenic factors such as change in soil use or deforestation.

The species with the most widespread distribution in the state were *P. teocote* and *P. pseudostrabus*. These were recorded in the greatest number of cells, and had the AOO of greatest extension (between 10,000 and 14,000 km²). *P. teocote* is widespread not only in Puebla, but throughout the country, occurring in 23 states (Contreras-Medina & Luna-Vega, 2007). In contrast, *A. hickelii* had an AOO of 108 km², and *P. greggii* and *P. oocarpa* less than 20 km², which makes these species rare in the state. Consequently, their records are very scarce, with only 2 localities recorded for these latter 2 pine species (Ramos-Dorantes, 2015). *A. hickelii* and *P. greggii* are endemic to Mexico (Farjon, 2010). The latter is restricted to the Sierra Madre Oriental (Contreras-Medina, 2004).

The species–accumulation curve predicts 16 Pinaceae taxa in the state of Puebla, which corresponds to the total number used to perform the analysis. Thus, the analysis indicates that there has been a satisfactory botanical exploration of the family in the state. Farjon (2010) reported other species as distributed in Puebla, namely *P. devoniana*, and *P. maximinoi*, and possibly

Table 3
List of variables selected with PCA for the potential distribution models.

Category	Variables
(a) Climatic variables (Hijmans, Cameron, Parra, Jones, & Jarvis, 2005; http://www.worldclim.org/)	bio1 = annual mean temperature bio4 = temperature seasonality (standard deviation × 100) bio5 = maximum temperature of warmest month bio9 = mean temperature of driest quarter bio11 = mean temperature of coldest quarter bio12 = annual precipitation bio16 = precipitation of wettest quarter bio18 = precipitation of warmest quarter PPH = precipitation of the humid months (May–October) TS = mean temperature of the dry months (November–April)
(b) Seasonality climatic variables (López-Mata et al., 2012)	
(c) Soil properties (Cruz-Cárdenas, López-Mata, Ortiz-Solorio, et al., 2014)	CE = electric conductivity CO = organic carbon K = potassium Mg = magnesium MO = organic material Na = sodium pH = hydrogen potential
(d) Topographic attributes (GTOPO web)	Aspect DAH = anisotropic heating ConvrgrIn = convergence index Runoff Slope TRI = terrain rugosity index TWI = topographic humidity index
(e) Normalized vegetation indices for 2009	IVNFEB = February normalized index IVNMAR = March normalized index IVNAPR = April normalized index IVNS = dry months of year normalized index

P. pringlei. Each of these is reported from 1 or more of the adjacent states of Mexico, Hidalgo, or Veracruz (Farjon, 2010; Martínez, 1992). Except for 1 cultivated specimen of *P. devoniana*, we have not located collections of these species from Puebla (Ramos-Dorantes, 2015).

The jackknife analysis did not identify any of the 28 variables used in modeling as making a particularly strong contribution for all the species of Pinaceae. Instead, each model used a particular combination of variables. Average annual temperature (Bio01) was the most important variable for many species. For 7 it accounted for the greatest contribution (Table 4). The combination of variables that contributed to each model was so specific that we decided to treat *P. pseudostrobus* and *P. pseudostrobus* var. *apulcensis* independently because in addition to morphological differences, the 2 taxa occur in different environments, with *P. pseudostrobus* var. *apulcensis* tending to occur in more semi-arid areas. The jackknife analysis showed that the informative variables differed between the 2 taxa. In *P. pseudostrobus* var. *pseudostrobus*, mean temperature of the dry months (TS) contributed most to modeling its distribution, whereas in *P. pseudostrobus* var. *apulcensis* the most important variable was mean temperature of coldest quarter (Bio11). Because the environmental conditions that influence the distribution of these taxa are different, their potential ranges are different, with *P. pseudostrobus* var. *apulcensis* more widespread in central areas of the state, and *P. pseudostrobus* var. *pseudostrobus* more widespread in the north and southeast (Fig. 4).

The results have a pattern generally similar to the known richness, except that in some sites more species are predicted than are currently recorded. The regions with the greatest richness are predicted to potentially contain 7–10 species, and areas with fewer species coincide with those of lesser known richness. The models predict suitable areas for Pinaceae where no records were

Table 4
Environmental variables that were informative in the modeling of potential distributions based on jackknife results.

Species	Bio1	CE	MO	DAH	IVNAPR	K	Mg	pH	Bio16	IVNMAR	Aspect	Bio5	Bio11	PPH	TS	Bio12	CO	Slope	IVNFEB	Runoff	TWI	Bio4	ConvrgrIn	Na	IVNS	TRI
<i>Abies hickelii</i>	*																									
<i>A. religiosa</i>	*																									
<i>Pinus ayacahuite</i>								*																		
<i>P. cembroides</i>	*																									
<i>P. chiapensis</i>							*																			
<i>P. douglasiana</i>							*																			
<i>P. greggii</i>															*											
<i>P. hartwegii</i>	*																									
<i>P. leiophylla</i>								*																		
<i>P. montezumae</i>	*																									
<i>P. oocarpa</i>		*																								
<i>P. patula</i>		*																								
<i>P. pseudostrobus</i>																										
<i>P. pseudostrobus</i> var. <i>apulcensis</i>													*													
<i>P. teocote</i>	*																									
<i>Pseudotsuga menziesii</i> var. <i>glauca</i>	*																									

The shaded cells indicate the variables that together contributed 85% of the importance per taxon.

* Variable with the greatest contribution in each taxon.




Figure 4. Potential distribution of *Pinus pseudostrabus* (left) and *P. pseudostrabus* var. *apulcensis* (right) in Puebla.

found, such as the border with the state of Veracruz, northeast of Pico de Orizaba, and the mountainous region west of Tehuacán. Botanical exploration in these areas is needed to improve our understanding of the patterns of distribution of the family in the state. The potential distribution area for species with fewer than 7 records, such as *A. hickelii*, for which the models included records from neighboring states, had small ranges (204 km²) within the state (Fig. 3). By contrast, taxa such as *P. pseudostrabus* var. *apulcensis*, with 46 records exclusive to Puebla, had a wide potential distributional area (9,897 km²) (Table 2).

The AOO of each species was only estimated for the state of Puebla. This regional focus offers relevant information for the conservation of these species that could be taken into account by local conservationists and authorities. Of all taxa, 4 had an AOO greater than 10,000 km² (*P. patula*, *P. pseudostrabus*, *P. pseudostrabus* var. *apulcensis*, and *P. teocote*), placing them in the category of Least Concern, whereas the remaining species were placed in a risk category. Six species are Near Threatened (*A. religiosa*, *P. ayacahuite*, *P. cembroides*, *P. hartwegii*, *P. leiophylla*, and *P. montezumae*), 2 species are in the category of Vulnerable (*P. chiapensis* and *Pseudotsuga menziesii* var. *glauca*), 2 species are Endangered (*A. hickelii* and *P. douglasiana*), and 2 species are Critically Endangered (*P. greggii* and *P. oocarpa*). Records for *P. greggii* and *P. oocarpa* in the state are scarce (Ramos-Dorantes, 2015) and their current AOO is very restricted (<5 km²). These regional categories agree in some cases with the IUCN Red List (IUCN, 2014) at the global level, as is the case of *P. greggii* and *A. hickelii*, because within the

country their distribution is restricted to only a few states (Farjon & Filer, 2013). By contrast, the evaluation of the regional level of risk for species in Puebla differs from the global evaluation in the cases of *P. oocarpa* and *P. douglasiana*, categorized as Least Concern because they have wide distributions at the national level (Farjon & Filer, 2013). One reason why *P. greggii* and *P. oocarpa* are categorized as Critically Endangered for Puebla is that in this study the only collections recorded were from montane cloud forest (Farjon, 2010; Ramos-Dorantes, 2015). This vegetation type is restricted in the state, making up approximately 2.1% of its area (Inegi, 2000). Fortunately, the models predict a larger potential AOO for these species with few records. For most of them the AOO is double, although more botanical exploration is needed to confirm their presence in these areas. Unfortunately, a substantial loss of natural vegetation is apparent in the soil use and vegetation maps (Inegi, 2000), which indicate that 78% of the state's area has been converted to crops, pastures, and urban areas.

Climate change and population growth are also expected to bring as a consequence the destruction of natural habitats and thus the potential distribution could diminish for all these species, not only for those with a restricted distribution. Conservative climate change scenarios predict a temperature increase of up to 1.7 °C, whereas the most severe scenarios predict up to 4.8 °C (IPCC, 2014). This, coupled with an increase in the human population, which could reach between 9 and 13.2 billion people by 2100 (Gerland et al., 2014), could bring losses in biodiversity. The various possible effects of climate change




Figure 5. Potential distribution of Pinaceae in Puebla ($N = 16$ taxa). The tones indicate the number of potential species.

may operate at different scales, from biomes, ecosystems, communities, species, populations or individuals, affecting their current distribution (Bellard, Bertelsmeier, Leadley, Thuiller, & Courchamp, 2012). An estimated 60–70% of temperate forests and other vegetation types in Mexico are predicted to be affected by climate change, with the tendency of species with temperate climate affinities to reduce their distribution (Villers-Ruíz & Trejo-Vázquez, 1998, 2004). As such, distribution models under climate change scenarios would be useful for predicting how the distribution of the family in the state would change. Conabio has recognized several terrestrial regions of priority (RTP's) for conservation within Puebla, such as the montane cloud forests of the Sierra Madre Oriental, Malinche, Cuetzalan, and Pico de Orizaba (Arriaga et al., 2000). These regions coincide with the areas of highest known species richness (Fig. 1) and potential richness (Fig. 5) of Pinaceae. Thus, conserving these regions would protect the species richness of the family in Puebla. However, it is also important to create individual conservation programs for the species with restricted distributions in the state.

Forests are the natural vegetation type of greatest coverage in Puebla (17.5%), and of these, conifer forests have the greatest extension, with 9.7% (Inegi, 2000). Although the status of other conifer families is unknown for the state, the present study suggests that botanical exploration of Pinaceae has been relatively satisfactory. It is possible that our knowledge of the taxa present in Puebla is nearly complete.

Acknowledgements

To Alejandro López Reyes, Sergio Hernández León, Patricia Rosas Escobar, Dylan Burge, and Rafael Torres Colín for their participation in the field. The first author thanks Conacyt (CB-2013, No. 221694) for providing an undergraduate thesis scholarship. To Conacyt (INFR-2016-01, No. 269382) for an infrastructure grant providing mobile cabinets for the specimens housed at MEXU.

References

- Aguirre-Gutiérrez, J., & Duivenvoorden, J. F. (2010). Can we expect to protect threatened species in protected areas? A case study of the genus *Pinus* in Mexico. *Revista Mexicana de Biodiversidad*, *81*, 875–882.
- Aguirre-Gutiérrez, J., Serna-Chávez, H. M., Villalobos-Arámula, A. R., Pérez-de la Rosa, J. A., & Raes, N. (2015). Similar but not equivalent: ecological niche comparison across closely-related Mexican white pines. *Diversity and Distributions*, *21*, 245–257.
- Arita-Watanabe, H. T. (2001). *Patrones geográficos de diversidad de los mamíferos terrestres de América del Norte*. Mexico City: Universidad Nacional Autónoma de México. Instituto de Ecología. Informe Final SNIB-Conabio proyecto No. Q068
- Arriaga, L., Espinoza, J. M., Aguilar, C., Martínez, E., Gómez, L., & Loa, E. (2000). *Regiones terrestres prioritarias de México*. Mexico City: Comisión Nacional para el Conocimiento y Uso de la Biodiversidad.
- Baldwin, R. A. (2009). Use of maximum entropy modeling in wildlife research. *Entropy*, *11*, 854–866.
- Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters*, *15*, 365–377.
- Colwell, R. K. (2009). *EstimateS, Version 8.2.0: Statistical estimation of species richness and shared species from samples (Software and User's Guide)*. Freeware published at: <http://viceroy.eeb.uconn.edu/estimates>
- Colwell, R. K., & Coddington, J. A. (1994). Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society (Series B)*, *345*, 101–118.
- Contreras-Medina, R. (2004). Gimnospermas. In I. Luna, J. J. Morrone, & D. Espinosa (Eds.), *Biodiversidad de la Sierra Madre Oriental* (pp. 137–148). Mexico City: Las prensas de Ciencias.
- Contreras-Medina, R., Castañeda-Aguado, D., & González-Zamora, A. (2007). Gimnospermas. In I. Luna, J. J. Morrone, & D. Espinosa (Eds.), *Biodiversidad de la Faja Volcánica Transmexicana* (pp. 129–138). Mexico City: Universidad Nacional Autónoma de México.
- Contreras-Medina, R., & Luna-Vega, I. (2007). Species richness, endemism and conservation of Mexican gymnosperms. *Biodiversity and Conservation*, *16*, 1803–1821.
- Cruz-Cárdenas, G., López-Mata, L., Ortiz-Solorio, C. A., Villaseñor, J. L., Ortiz, E., Silva, J. T., et al. (2014). Interpolation of Mexican soil properties at a scale of 1:1,000,000. *Geoderma*, *213*, 29–35.
- Cruz-Cárdenas, G., López-Mata, L., Villaseñor, J. L., & Ortiz, E. (2014). Potential species distribution modeling and the use of principal component analysis as predictor variables. *Revista Mexicana de Biodiversidad*, *85*, 189–199.
- Cruz-Cárdenas, G., Villaseñor, J. L., López-Mata, L., & Ortiz, E. (2013). Distribución espacial de la riqueza de especies de plantas vasculares en México. *Revista Mexicana de Biodiversidad*, *84*, 1189–1199.
- Delgadillo, C., Villaseñor, J. L., Cárdenas, Á., & Ortiz, E. (2014). Diversity and distribution of mosses in the state of Hidalgo, Mexico. *Revista Mexicana de Biodiversidad*, *85*, 84–97.
- Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., et al. (2013). Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography*, *36*, 27–46.
- Eckenwalder, J. E. (2009). *Conifers of the world*. Portland, Oregon: Timber Press.

- Evangelista, P. H., Kumar, S., Stohlgren, T. J., & Young, N. E. (2011). Assessing forest vulnerability and the potential distribution of pine beetles under current and future climate scenarios in the Interior West of the US. *Forest Ecology and Management*, 262, 307–316.
- Evans, J. S., & Cushman, S. A. (2009). Gradient modeling of conifer species using random forests. *Landscape Ecology*, 24, 673–683.
- Farjon, A. (2010). *A handbook of the world's conifers*. Leiden, Netherlands: Brill.
- Farjon, A., & Filer, D. (2013). *An atlas of the world's conifers: an analysis of their distribution, biogeography, diversity and conservation status*. Leiden, Netherlands: Brill Academic Publishers.
- Farjon, A., & Styles, B. T. (1997). *Pinus (Pinaceae)*. *Flora neotropica monograph* 75. New York: The New York Botanical Garden.
- García-Aranda, M. A., Cantú-Ayala, C., Estrada-Castillón, E., Pando-Moreno, M., & Moreno-Talamantes, A. (2012). Distribución actual y potencial de *Taxus globosa* (Taxaceae) en México. *Journal of the Botanical Research Institute of Texas*, 6, 587–598.
- GBIF (Global Biodiversity Information Facility Secretariat). (2014). *GBIF backbone taxonomy*. [accessed 14 Dec 2016]. Retrieved from: <http://www.gbif.org/species/3925>
- Gerland, P., Raftery, A. E., Ševčíková, H., Li, N., Gu, D., Spoorenberg, T., et al. (2014). World population stabilization unlikely this century. *Science*, 346, 234–237.
- Gernandt, D. S., & Pérez-de la Rosa, J. A. (2014). Biodiversidad de Pinophyta (coníferas) en México. *Revista Mexicana de Biodiversidad*, 85(Suppl.), S126–S133.
- Gómez-Mendoza, L., & Arriaga, L. (2007). Modeling the effect of climate change on the distribution of oak and pine species of Mexico. *Conservation Biology*, 21, 1545–1555.
- Gotelli, N. J., & Colwell, R. K. (2001). Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecological Letters*, 4, 379–391.
- Graham, C. H., & Hijmans, R. J. (2006). A comparison of methods for mapping species ranges and species richness. *Global Ecology and Biogeography*, 15, 578–587.
- Gugger, P. F., González-Rodríguez, A., Rodríguez-Correa, H., Sugita, S., & Cavender-Bares, J. (2011). Southward Pleistocene migration of Douglas-fir into Mexico: phylogeography, ecological niche modeling, and conservation of 'rear edge' populations. *New Phytologist*, 189, 1185–1199.
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25, 1965–1978.
- Inegi (Instituto Nacional de Estadística Geografía e Informática). (2000). *Síntesis geográfica del estado de Puebla*. México [CD-ROM].
- Inegi (Instituto Nacional de Estadística Geografía e Informática). (2004). *Las provincias fisiográficas de México y sus subdivisiones. Sistema fisiográfico DGGTENAL*. Mexico City: Escrito de la Subdirección de actualización de Recursos Naturales.
- IPCC (Intergovernmental Panel on Climate Change). (2014). In Core Writing Team, R. K. Pachauri, & L. A. Meyer (Eds.), *Climate change 2014: synthesis report. Contribution of working groups I, II and III to the fifth assessment report of the intergovernmental panel on climate change*. Geneva, Switzerland: IPCC.
- IUCN. (2001). *IUCN red list categories and criteria: Version 3.1*. Gland, Switzerland and Cambridge, UK: IUCN Species Survival Commission, IUCN.
- IUCN. (2014). *The IUCN red list of threatened species. Version 2014.2*. <http://www.iucnredlist.org>
- Janžekovič, F., & Novak, T. (2012). PCA – A powerful method for analyze ecological niches. In P. Sanguansat (Ed.), *Principal component analysis – multidisciplinary applications* (pp. 127–142). Rijeka, Croatia: InTech.
- Jiménez-Valverde, A., & Hortal, J. (2003). Las curvas de acumulación de especies y la necesidad de evaluar la calidad de los inventarios biológicos. *Revista Ibérica de Aracnología*, 8, 151–161.
- Leal-Nares, Ó., Mendoza, M. E., Pérez-Salicip, D., Geneletti, D., López-Granados, E., & Carranza, E. (2012). Distribución potencial del *Pinus martinicensis*: un modelo espacial basado en conocimiento ecológico y análisis multicriterio. *Revista Mexicana de Biodiversidad*, 83, 1152–1170.
- López-Mata, L., Villaseñor, J. L., Cruz-Cárdenas, G., Ortiz, E., & Ortiz-Solorio, C. (2012). Predictores ambientales de la riqueza de especies de plantas del bosque húmedo de montaña de México. *Botanical Sciences*, 90, 27–36.
- López-Reyes, A., Pérez-de la Rosa, J. A., Ortiz, E., & Gernandt, D. S. (2015). Morphological, molecular, and ecological divergence in *Pinus douglasiana* and *P. maximinoi*. *Systematic Botany*, 40, 658–670.
- Martínez, M. (1992). *Los pinos mexicanos*. Mexico City: Ediciones Botas.
- Moat, J. (2007). *Conservation assessment tools, extension for ArcView 3.x, version 1.2. GIS Unit*. Kew, United Kingdom: Royal Botanic Gardens.
- Ogawa-Onishi, Y., Berry, P. M., & Tanaka, N. (2010). Assessing the potential impacts of climate change and their conservation implications in Japan: a case study of conifers. *Biological Conservation*, 143, 1728–1736.
- Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., et al. (2011). *Ecological niches and geographic distributions*. Princeton, NJ: Princeton University Press.
- Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190, 231–259.
- Ramos-Dorantes, D. B. (2015). *Revisión taxonómica de la familia Pinaceae en el estado de Puebla* Undergraduate Thesis. Puebla: Escuela de Biología, Benemérita Universidad Autónoma de Puebla.
- R Development Core Team. (2015). *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. ISBN 3-900051-07-0. Available at: <http://www.R-project.org>
- Rzedowski, J. (1978). *Vegetación de México*. Mexico City: Limusa.
- Sánchez, G. A. (2008). Una visión actual de la diversidad y distribución de los pinos de México. *Madera y Bosques*, 14, 107–120.
- (2014). *SEInet (Southwest Environmental Information Network)*. <http://swbiodiversity.org/seinet/collections/index.php>
- Soberón, J. M., & Llorente, J. B. (1993). The use of species accumulation functions for the prediction of species richness. *Conservation Biology*, 7, 480–488.
- Soberón, J., & Peterson, A. T. (2005). Interpretation of models of fundamental ecological niches and species' distributional areas. *Biodiversity Informatics*, 2, 1–10.
- StatSoft, Inc. (2011). *STATISTICA (data analysis software system), version 10*. Available at: <http://www.statsoft.com>
- Suárez-Mota, M. E., & Villaseñor, J. L. (2011). Las compuestas endémicas de Oaxaca, México: diversidad y distribución. *Boletín de la Sociedad Botánica de México*, 88, 55–66.
- The New York Botanical Garden: Herbarium of The New York Botanical Garden (NY) (2014) [accessed 16 Dec 2014]. Retrieved from: <http://www.gbif.org/dataset/7133ff0a-f762-11e1-a439-00145eb45e9a>
- The United States National Herbarium (US) (2014) [accessed 16 Dec 2014]. Retrieved from: http://botany.si.edu/colls/Collections_history.htm
- Villaseñor, J. L., Ortiz, E., Cadena, J., & Estrada, A. S. (2013). Patrones de riqueza florística en el estado de Jalisco: la tribu Senecioneae (Asteraceae) como estudio de caso. *Ibugana*, 4, 63–78.
- Villers-Ruiz, L., & Trejo-Vázquez, I. (1998). Climate change on Mexican forests and natural protected areas. *Global Environmental Change*, 8, 141–157.
- Villers-Ruiz, L., & Trejo-Vázquez, I. (2004). Evaluación de la vulnerabilidad en los ecosistemas forestales. In J. Martínez, & A. Fernández (Eds.), *Cambio climático: una visión desde México* (pp. 239–254). Mexico City: Secretaría de Medio Ambiente.