

Taxonomy and systematics

## ***Hymenolepis ackerti* n. sp. (Eucestoda: Hymenolepididae) infecting cricetid rodents from the central Great Plains of North America**

### ***Hymenolepis ackerti* n. sp. (Eucestoda: Hymenolepididae) de roedores cricétidos recolectados en las planicies centrales de Norteamérica**

Brian Rowan <sup>a</sup>, Andrew G. Hope <sup>b</sup>, F. Agustín Jiménez <sup>a, \*</sup>

<sup>a</sup> Southern Illinois University, School of Biological Sciences, 1125 Lincoln Drive, 355C Life Science 2, School of Biological Sciences, Southern Illinois University, Carbondale Illinois 62901-6501, USA

<sup>b</sup> Kansas State University, Division of Biology, 116 Ackert Hall, Division of Biology, Kansas State University, Manhattan, KS 66506 USA

\*Corresponding author: agustinjz@zoology.siu.edu (F.A. Jiménez)

Received: 13 January 2022; accepted: 28 November 2022

<http://zoobank.org/urn:lsid:zoobank.org:pub:E74CB8AA-FF75-49FD-97D1-B8B0519C7E5D>

#### **Abstract**

*Hymenolepis ackerti* n. sp., parasite of rodents from the tallgrass prairie ecoregion of North America is herein characterized. This tapeworm occurs in 3 species of rodents including the hispid cotton rat *Sigmodon hispidus*, the eastern woodrat *Neotoma floridana*, and the prairie vole *Microtus ochrogaster*. A comparison against the other 10 congeneric species known from North America reveals that this species is different based on the size of the scolex, length of rostellar capsule, testicular arrangement, and the size of cirrus sac, seminal receptacle and eggs. A comparison of mitochondrial DNA reveals that tapeworms present in sympatric mammals share the same mitochondrial haplotype and feature similar morphology, supporting their recognition as a single species. The phylogenetic position of *H. ackerti* relative to other species is still to be resolved, since there are no homologous sequences available for most species in the genus. Given the pervasiveness of these parasites across rodents in the continent, we recommend diligence among scientists to build public archives of tapeworm specimens collected from mammals across North America, and globally. In the present manuscript, we propose a method to sample DNA while still allowing specimens to be postfixed for staining or fluid-preserved for long term storage.

**Keywords:** Konza Prairie; Tallgrass prairie; Expected identification; Host unspecificity

## Resumen

Se presenta la caracterización de *Hymenolepis ackerti* esp. n., parásito de roedores de las praderas centrales norteamericanas. Este céstodo infecta 3 especies simpátricas de roedores *Sigmodon hispidus* (Sigmodontinae), *Neotoma floridana* (Neotominae) y *Microtus ochrogaster* (Arvicolinae). La comparación directa con las 10 especies congénéricas presentes en Norteamérica revela diferencias en el tamaño del escólex, la longitud de la cápsula rostellar, el arreglo testicular y el tamaño del saco del cirro, receptáculo seminal y huevos. La comparación de haplotipos mitocondriales y la similitud de los caracteres morfológicos indican la conespecificidad de céstodos que infectan roedores de estas 3 especies simpátricas. La posición de la especie dentro de *Hymenolepis* es aún incierta, ya que se carece de secuencias disponibles para la mayoría de las especies del género que posibiliten su reconstrucción filogenética. Dada la ubiquidad de roedores y sus parásitos, recomendamos que científicos practicantes generen un acervo de céstodos presentes en roedores tanto en Norteamérica como del mundo. En el presente trabajo, proponemos un método para muestrear ADN que permite la posfijación de los parásitos, misma que favorece su tinción o incluso su almacenamiento permanente en un medio fluido.

*Palabras clave:* Pradera Konza; Pradera pastizales altos; Identificación esperada; Inespecificidad hospedatoria

## Introduction

*Hymenolepis* Weinland, 1858 includes species that feature an unarmed scolex with a rudimentary rostellar apparatus, anastomoses connecting ventral excretory canals, saccate uterus and testes arranged triangularly or in a transversal line. There are 10 species known to occur in North America which infect sciurid, arvicoline, geomyid and neotomine rodents (Gardner et al., 2020; Hoberg et al., 2016; Makarikov et al., 2015). In addition, the cosmopolitan *H. diminuta* Weinland, 1858 has been recorded throughout the continent infecting humans and both synanthropic and wild rodents (Coggins & McDaniel, 1975; Faulkner & Lochmiller, 2000; Harkema, 1946; Harkema & Kartman, 1948; Mollhagan, 1978; Seidenberg et al., 1974).

Although the majority of *Hymenolepis* species are known to infect only their definitive host, two species occur in unrelated sympatric mammals. These include *H. citelli* (McLeod, 1933) and *H. folkertsi* Makarikov, Nims, Galbreath and Hoberg, 2015; the latter was documented from a variety of unrelated species of cricetid, geomyid, heteromyid and sciurid rodents distributed across the United States (Gardner, 1985; Gardner et al., 2020; Hoberg et al., 2016; Makarikov et al., 2015; Pfaffenberger et al., 1985; Riley & Shannon, 1922; Smith et al., 1953). Recently, the use of mitochondrial genes has assisted in establishing the species identity of individual tapeworms identified as *H. folkertsi* collected across the vast geography of North America and from a wide diversity of mammals. By analyzing sequence divergence, Hoberg et al. (2016), established inter-population genetic structure and related this to the characterization of intraspecific morphological variability of diagnostic characters. They also determined that the genetic distance among individuals of *H. folkertsi*

across the distribution of the species oscillates between 1.9 and 2.7% (Hoberg et al., 2016). A similar approach has been used to evaluate the intraspecific genetic diversity in species of the hymenolepidid *Arostrilepis* Mas-Coma and Tenora, 1997 (Makarikov et al., 2020) and provide identifications for human-dwelling hymenolepidids (Nkouawa et al., 2016).

Ongoing biodiversity monitoring of small mammals and their associated ecto- and endo-parasite faunas on the Konza Prairie Biological Station (KPBS), and Long-Term Ecological Research (LTER) site, located in northeastern Kansas, USA, reflects 6 consecutive years of both mark-recapture data and comprehensive mammal and parasite voucher collection (Galbreath et al., 2019; Hope, 2019). These integrated surveys are performed with the purpose of documenting the distributional ecology, population biology and co-evolutionary history of these communities, as they occur across a landscape scale experimental manipulation of land-use treatments. These treatments include a factorial design of controlled wildfire and grazing regimes that have been continuously managed for > 40 years of ecological experimentation focused on understanding drivers of grassland resilience. The KPBS is located in the Flint Hills ecoregion and supports the largest remaining tract of native tallgrass prairie through the central Great Plains. Experimental manipulation, especially through wildfire suppression has resulted in a mosaic landscape of native prairie, shrubland communities, and late-seral woodlands. Small mammals were sampled in multiple habitats reflected by variable fire return intervals. Although most of this site is located on terraced limestone and has never been suitable for agriculture, the regional impacts of changing land use has profoundly altered most tallgrass prairie habitats (Briggs et al., 2005), associated mammalian community structure (Bruckerhoff et al.,

2021), and ecosystem function (Ratajczak et al., 2016). To date, the lack of a detailed species level inventory of parasites in this locality precludes any knowledge of their biological role in this ecosystem.

Here we report some findings from our efforts to identify the species of tapeworms present in tallgrass prairie rodents and propose the description of a new species recovered from hispid cotton rats *Sigmodon hispidus* Say and Ord, 1825, eastern woodrats *Neotoma floridana* (Ord, 1818), and prairie voles *Microtus ochrogaster* (Wagner, 1842). Our description is based on the examination of the morphological features that characterize these parasites as well as their molecular identity based on mitochondrial genes.

## Materials and methods

Mammals were trapped using Sherman® live traps (H.B. Sherman, Tallahassee, FL). Subsequently they were euthanized under approved protocols (Hope IACUC #3579, #4055) and processed to preserve all specimen parts (Galbreath et al., 2019) for permanent archive in the Museum of Southwestern Biology, University of New Mexico (<http://arctos.database.museum/>). The trapping design used at the Konza Prairie Biological Station (KPBS), and Long-Term Ecological Research (LTER) site, (Kansas, USA, 39.07264° N, 96.57818° W, 416 m elevation) is detailed in the data report Universally available (Hope et al., 2019). For parasite preservation, the digestive system was cut open and the tapeworms present were removed. Worms were washed and killed with hot water to induce their relaxation. Specimens were preserved in 70% ethanol, washed, postfixed in 4% formaldehyde and stained in Semichon's paracarmine to be mounted permanently in Canada Balsam. Specimens identified as *H. diminuta* collected from hispid cotton rats from Oklahoma were borrowed from the National Parasite Collection (USNM1382519 = USNPC87319) and used for comparison. All measurements are in micrometers (µm), unless otherwise noted. Type material was deposited at Museum of Southwestern Biology (MSB), Division of Parasites, University of New Mexico, Albuquerque, New Mexico, USA; Harold W. Manter Laboratory of Parasitology (HWML), University of Nebraska, Lincoln, Nebraska, USA, and Colección Nacional de Helminthos (CNHE), Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, Mexico.

Prior to the staining and mounting of specimens, fragments of a few contiguous mature proglottids were excised and used for the extraction of genomic DNA using a commercial kit (Qiagen, Valencia CA), whereas the rest of the specimen was processed as described above for

morphological examination. The mitochondrial NADH dehydrogenase subunit 1 (Nad1) gene was amplified using primers and thermal profiles developed and described elsewhere (Haukisalmi et al., 2010; Morgan & Blair, 1998). The resulting amplicons were cleaned using ExoSAP (GE Healthcare, Cleveland, OH) and submitted to a commercial facility to be sequenced using Sanger chemistry (Eurofins Inc., Lexington, Kentucky). Sequences were assembled in Sequencher 3.5 (Gene Codes Corp. Ann Arbor, Michigan), and aligned and trimmed with available homologous sequences from hymenolepidids and its putative related groups *Arostrilepis*, *Staphylocystis* Villot, 1877 and *Staphylocystoides* Yamaguti, 1959 using Clustal W (Sievers et al., 2011) with default settings. Accession numbers of the sequences used are related in Table 1. Resulting alignments had a size of 712 bp. Following alignment, the model of evolution GTR was selected using the best fit model according to the corrected Akaike Information Criterion as implemented in JModeltest (Posada, 2008). The posterior probability of branches was calculated using Bayesian inference as implemented in Mr. Bayes v3.2.7 using the CIPRES portal (Miller et al., 2010). Trees were visualized using FigTree v1.4.3 (Rambaut, 2012). Genetic distances were estimated with PAUP\*.

## Description

Genus *Hymenolepis* Weinland, 1858

*Hymenolepis ackerti* n. sp. (Figs. 1, 2)

<http://zoobank.org/urn:lsid:zoobank.org:act:170E75EE-96A9-40B9-B814-3A54CF26D6BE>

Based on 10 specimens, measurements of the holotype in parenthesis. Scolex unarmed, semispherical, 417-570 (548) long by 438-567 (472) wide (Fig. 1A). Rostellar capsule 100-138 (100) long by 66-87 (77) wide, thin walls not reaching the posterior margin of suckers; anterior canal or apical organ 31-36 (31). Osmoregulatory canals terminate in scolex, near posterior margin of rostellar capsule (Fig. 1A). Suckers unarmed (n = 40), slightly oval, 112-192 (126 - 169) long by 103-182 (145-160) wide, endowed with thick muscular walls. Neck 293-423 (423) long, conspicuously narrower than scolex (Fig. 1A). Relatively small fully developed strobila, length 40.4-106.6 mm (106.6, n = 8), maximum strobilar width 1,084-2,207, (2,207 mm) at level of newly gravid proglottids. Ventral excretory canals 62-93 (93) wide, connected with transversal anastomoses. Dorsal excretory canal 7-10 (9) wide. Development of proglottids protandrous; proglottids feature non-alternating dextral genital pores (Fig. 1B).

Proglottids anapolytic wider than long; strobilar margins craspedote, with conspicuous intersegmental divisions between mature and gravid proglottids. Mature

Table 1

List of specimens and sequences used to analyze the phylogenetic position of *Hymenolepis ackerti*.

Species	ND1 GenBank No.	Collection number	Locality	Host species
<i>Arostrilepis beringiensis</i>	KM516216		Irkutsk, Russia	<i>Lemmus sibiricus</i>
<i>Hymenolepididae</i>	JQ950694	MHNGE-INVE 79505	Hualaihue, Chile	<i>Sephanoides sephaniodes</i>
<i>Hymenolepis ackerti</i>	OP649625	MSB: Para: 35315	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649626	HWML118076	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649633	MSB: Para: 35316	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649634	HWML118077	Kansas, USA	<i>Neotoma floridana</i>
<i>Hymenolepis ackerti</i>	OP649632	MSB: Para: 35317	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649631	MSB: Para: 35318	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649630	MSB: Para: 35319	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649629	MSB: Para: 35314	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649628	MSB: Para: 35313	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis ackerti</i>	OP649627	HWML118078	Kansas, USA	<i>Sigmodon hispidus</i>
<i>Hymenolepis diminuta</i>	AF314223	N/A	Lab, Michigan	Lab isolate
<i>Hymenolepis diminuta</i>	AP017664	N/A	Lab, Japan	Lab isolate
<i>Hymenolepis diminuta</i>	LR536429	N/A	Lab strain WMS-il1, Poland	Lab Isolate
<i>Hymenolepis diminuta</i>	HM149290	N/A	Madagascar	<i>Rattus rattus</i>
<i>Hymenolepis diminuta</i>	HM149291	N/A	Canary Islands, Spain	<i>Rattus rattus</i>
<i>Hymenolepis folkertsi</i>	OP649624	35314	Kansas, USA	<i>Peromyscus leucopus</i>
<i>Hymenolepis folkertsi</i>	KX782315	N/A	Pennsylvania, USA	<i>Peromyscus leucopus</i>
<i>Hymenolepis folkertsi</i>	KX792199	MSB: Para: 23512	Idaho, USA	<i>Tamias amoenus</i>
<i>Hymenolepis folkertsi</i>	KX792195	MSB: Para: 23503	Michigan, USA	<i>Peromyscus leucopus</i>
<i>Hymenolepis nana</i>	AP017666	N/A	Lab, Japan	Lab isolate
<i>Hymenolepis</i> sp. C31	HM149299	N/A	Madagascar	<i>Eliurus tanala</i>
<i>Hymenolepis</i> sp.	HM149298	N/A	Madagascar	<i>Tenrec ecaudatus</i>
<i>Hymenolepis</i> sp. U45	HM149292	N/A	Kazakhstan	<i>Microtus socialis</i>
<i>Hymenolepis</i> sp. AY0	HM149295	N/A	Wyoming, USA	<i>Thomomys</i> sp.
<i>Hymenolepis</i> sp. BT0	HM149296	N/A	Victoria, Australia	<i>Rattus fuscipes</i>
<i>Hymenolepis</i> sp. CA9	HQ589353	N/A	South Korea	<i>Apodemus agrarius</i>
<i>Hymenolepis</i> sp. U9	HM149297	N/A	Turkey	<i>Apodemus sylvaticus</i>
<i>Hymenolepis</i> sp. Z39	HM149289	N/A	Bosnia and Herzegovina	<i>Microtus subterraneus</i>

proglottids 97-225 (175-225, n = 64) long by 1,260-1,771 (1,587-1,771, n = 64) wide. Genital atrium simple, infundibular, located at anterior end of proglottid margin 18-29 (18). Cirrus sac pyriform and muscular, relatively long 142-266 (142-256, n = 128) by 32-61 (32-61, n = 82); antiporal portion of cirrus sac occasionally reaches ventral excretory canal, yet never crosses longitudinal osmoregulatory canals (Fig. 1B, C). Cirrus claviform, armed with minute spines 73-101 (78-97, n = 8). Internal

seminal vesicle oval, 73-186 (73-166, n = 42) more than half of cirrus sac length (Fig. 1B, C); external seminal vesicle fusiform 166-235 (205-273, n = 36), dorsoventrally undulating and distinctly shorter than seminal receptacle (Fig. 1B, C). Genital ducts cross dorsally both ventral and dorsal osmoregulatory canals. Usually 3 testes, 1 poral, 2 antiporal; 4 or 5 testes rarely occur. Testes oval, somewhat linearly arranged 88-205 (147-205, n = 120) long by 123-273 (161-245, n = 120) wide; poral testis separated from




Figure 1. *Hymenolepis ackerti* n. sp. A, Scolex showing anterior portion of excretory canals that do not anastomose or reach the anterior canal. Scale bar = 100  $\mu$ m; B, detail of genitalia showing genital atrium, cirrus sac containing both cirrus and internal seminal vesicle, and distal end of the external seminal vesicle. The connection of the vagina with the seminal receptacle is shown in the lower proglottid. Scale bar = 100  $\mu$ m; C, dorsal view of a mature proglottid showing the relative position of the ovary, seminal receptacle, location of testes, sigmoidal external seminal receptacle, and cirrus sac. Scale bar = 200  $\mu$ m; D, uterine egg featuring hooks in onchosphere. Scale bar = 25  $\mu$ m.

antiporals by female genitalia (Fig. 1C). Vaginal opening ventral to cirrus opening, copulatory part of vagina 75-107 (75-88, n = 16) long (Fig. 1B). Fusiform seminal receptacle, 520-714 (608-710, n = 24) (Fig. 1C). Ovary median, fan-shaped, 45-159 (85-106, n = 46) long by 108-304 (169-188, n = 41) wide, occasionally overlapping proximal poral testis. Vitellarium lobed, postovarian 50-169 (80-169, n = 40) long by 72-200 (72-143, n = 40) wide.

Uterus develops laterally from the central part of the proglottid and elongates to fill the space between the excretory canals and the space closer to lateral margins (Fig. 2A, B); saccular projections dorsally directed. In gravid proglottids the saccular uterus fills the entirety of the proglottid, eggs occupy first the space between excretory canals and subsequently eggs fill the marginal spaces in terminal proglottids (Fig. 2B). Gravid proglottids 275-803 (548-803, n = 76) long by 1,084 - 2,207 (1,851-2,207, n = 76) wide.

Outer coat of eggs usually spherical 38-55 (35-36, n = 50) long by 38-56 (40-45, n = 50) wide, surface endowed with uniform embossment; onchosphere 18-33 (20-33)

long by 19-37 (22-37) wide. Embryophore 25-36 (25-29) long by 34-40 (34-35) wide (Fig. 1D). Pair of slender median hooks, 8-13 (9-12, n = 15); anterolateral hooks relatively robust 11-16 (11-15, n = 43). Terminal or last proglottids contain no eggs (Fig. 2D).

#### Taxonomic summary

*Symbiotype*: hispid cotton rat, *Sigmodon hispidus* (Say and Ord).

*MSB*: Mamm: 330681.

*Site of Infection*: small intestine.

*Type locality*: USA, Kansas, Konza Prairie Biological Station (KPBS), and Long-Term Ecological Research (LTER) site, 39°4'22" N, 96°34'41" W, 416 m elevation.

*Specimens deposited*: holotype (MSB: Para: 35308). Paratypes MSB: Para: 35309, HWML216883, HWML118076, CNHE11734. Vouchers: MSB: Para: 35306-35307, 35310-35311; 35313-35319; HWML118077 and 118078, CNHE11735.

*Other hosts*: prairie vole, *Microtus ochrogaster* (Wagner) and the eastern woodrat, *Neotoma floridana* (Ord)




Figure 2. Micrographs of uterine development of *Hymenolepis ackerti* n. sp.(holotype). A, Ventral view of proglottids featuring developed uterine branches in the space between excretory ducts, scale bar = 500 µm; B, ventral view of terminal gravid proglottids, showing all uterine branches filled with eggs, scale bar = 1 mm.

*Specimens examined:* *Neotoma floridana* MSB: Mamm: 330678, *Microtus ochrogaster* MSB: Mamm: 331573 and *Sigmodon hispidus* MSB: Mamm: 305504, MSB: Mamm: 305616, MSB: Mamm: 330605, MSB: Mamm: 330663, MSB: Mamm: 331560, MSB: Mamm: 331542, MSB: Mamm: 331563, MSB: Mamm: 331584, MSB: Mamm: 331590, MSB: Mamm: 332155.

*NCBI GenBank numbers:* OP649625 - OP649634, OQ407494 - OQ407496

*Etymology:* the species is named after J. E. Ackert, former president of the American Society of Parasitologists and head of the former Zoology Department at Kansas State University.

#### Remarks

*Hymenolepis ackerti* is unique among all described species of this genus present in North America because of the extremely large and semispherical scolex, the small embryo hooks and the combination of a relatively large

cirrus, relatively large seminal receptacle, small eggs and the disposition of the rostellar capsule that barely reaches the anterior margin of the suckers. The rostellar capsule of *H. ackerti* is completely embedded in the scolex, which allows the differentiation of this species from *H. folkertsi*, *H. pitymi* Yarinsky, 1952, and *H. tualatinensis* Gardner, 1985; further, all 3 of these species feature testes arranged in a triangle, as opposed to the linear arrangement characteristic of *H. ackerti*.

In some individuals the seminal receptacle of *H. ackerti*, is relatively elongated and reaches the center of the proglottid. This is notable because this proportion is seen in only four other species present in the New World, which include *H. cratogeomys*, *H. diminuta*, *H. scalopi*, and *H. weldensis* Gardner and Schmidt, 1988. However, the eggs and hooks present in the embryo of *H. ackerti* are typically smaller than the homologous structures in those four species. In particular, *H. ackerti* shares six characteristics with *H. weldensis*, namely the configuration




Figure 3. Phylogenetic reconstruction showing the position of *Hymenolepis ackerti* relative to available sequences of the mtDNA ND1 locus for hymenolepidids. The values atop each node represent posterior probabilities (Bayesian inference) and values below represent Bootstrap support based on 1,000 replicates.

of the rostellar capsule, the testicular arrangement, the size of ovaries, length of seminal receptacle, size of vitelline glands and size and extension of cirrus sac. However, both species can be differentiated by the size of the eggs, onchosphere and embryonic hooks, being larger in *H. weldensis*.

Compared to *H. robertrauschi*, individuals of *H. ackerti* feature relatively smaller eggs and embryos, yet rostellar capsule and seminal receptacle are at least twice as long in *H. ackerti*. The larger size of the scolex and the smaller hooks in the onchosphere can be used to differentiate *H. ackerti* from the rest of the North American species. The scolex of *H. ackerti* is almost twice as wide as the homologous structure of *H. citelli*, *H. cratogeomys*, *H. gomydis* and *H. scalopi* (Table 2). Finally, only the eggs of *H. pitymi* appear to be smaller than those in *H. ackerti*, yet both species can be differentiated based on the arrangement of the testes, linear in the latter and triangular in the former. The relative size of the hooks in the embryo are similar to those present in *H. diminuta*; however, the testes and the eggs of *H. ackerti* are slightly larger, whereas the cirrus sac is usually longer in *H. diminuta*.

Genetic screening of individuals collected from different species of mammals

The monophyly of *H. ackerti* is supported by their phylogenetic placement according to the analyses of *ND1* mtDNA locus herein analyzed (Fig. 3). In the resulting

topologies, *H. ackerti* is not the sister species for either *H. diminuta* nor for *H. folkertsi*, the other species present in North America for which comparative GenBank sequences were available. All sequenced specimens of *H. ackerti* differ by one or two single point mutation representing three haplotypes of *ND1*, their intraspecific genetic distance oscillates between 0.15 and 0.3% (Table 3).

## Discussion

Cestode specimens collected from hispid cotton rats in the last 75 years were usually identified as *H. diminuta* (Coggins & McDaniel, 1975; Faulkner & Lochmiller, 2000; Harkema, 1946; Harkema & Kartman, 1948; Mollhagan, 1978; Seidenberg et al., 1974). From these, only few specimens collected from Oklahoma were deposited in scientific collections as verifiable voucher specimens. Examination of these specimens, USNM1392519 = USNPC87319, revealed they cannot be diagnosed as *H. diminuta*, based on the fact that they feature an everted armed rostellum, not typical for the species.

Relative to the intraspecific morphological variability, it must be noted that specimens of *H. ackerti* feature relatively short proglottids and most organs are arranged in the same dorsoventral plane. A similar situation occurs with the position of the vitellaria relative to the ovary. Further, the length of the strobila of some specimens

Table 2

Comparative measurements of relevant characters used in the taxonomy of *Hymenolepis* species from North America.

	<i>H. ackerti</i> n. sp.	<i>H. diminuta</i> (Rudolphi, 1819)	<i>H. citelli</i> (McLeod, 1933)	<i>H. crateogeomyos</i> Gardner, Dursahinan, Campbell and Rácz, 2020	<i>H. tualatinensis</i> Gardner, 1985	<i>H. weldensis</i> Gardner and Schmidt, 1988	<i>H. geomydis</i> Gardner and Schmidt, 1988	<i>H. folkertsi</i> Makarikov, Nims, Galbreath and Hoberg 2015	<i>H. scalopi</i> Schultz, 1939	<i>H. pitymi</i> Yarinsky 1952	<i>H. robertrauschi</i> Gardner, Luedders and Duszinski, 2014
Scolex with rostrum-like projection	No	No	No	No	Yes	No	No	Yes	No	Yes	No
Strobila length (mm)	106.57 (40-51)	183	150	88-129	24-210	111-165	72-168	99-116	100-200	19.6	42.4 - 83.4
Strobila width (mm)	1.52 (1.69-1.94)	2.38-2.56	2.8	2.72-3.41	1.75	0.82-0.94	1.98-3.3	1.71-1.85	2.57	750	1.18 - 2.54
Scolex width	472 (438-568)	286-296	245	209-227	92-167	126-288	194-245	168	170 - 232		199 - 257
Sucker size	126 - 169 × 145 - 160 (111-184 × 121-182)	96-108	113 × 87	84-89 × 109-117-	-	-	92-124 × 65-94	93-102 × 70-86	35-62 × 86-92		119 - 164 × 82 - 95
Rostellar pouch size (length × width)	100 × 77 (115- 138X66-87)	94-108 × 44-52		55-59	51-61 long	-	-	67 × 43 long	121		50 long
Mature proglottid size	-	-	-	-	-	-	-	90-150 × 810-1030			
Testes size (length × width)	93-200 × 131-272	118-220 × 40-48	143 × 113	141-190 × 95-121	63-141 × 54-141	92-166	55-180 × 81-180	75-112 × 44-62	101-123 × 87-113		
Testicular arrangement	Linear	Linear	Linear	Linear	Triangular	Linear	Triangular	Triangular	Linear	Triangular	Triangular
Cirrus-sac size (length × width)	143-256 × 32-61	240-300 × 28-48	157 long	121-145 × 45-57	56-150 × 26-49	149-194 × 34-51	80-160 × 36-67	138-154 × 30-39	125-159 × 42-49	79 long	147 - 233 × 33 - 61
Cirrus size	78-97 (72-102)	38-47*	-	—	-	-	-	45-58 × 7-10			84 - 157
Cirrus armature	minute spines	minute spines	-	armed	minute spines	minute spines	minute spines	minute spines	minute spines		
Ovary width	169-188 (108-246)		-	168-203	96-216	90-293	180-494	147-217	193-220	97-116	130 - 297
Vitellarium size (length × width)	80-169(50-89 × 87-167)	-	-	45-53X62-80	34-109 × 37-132	50-112 × 54-106	61-137 × 101-209	30-57 × 55-98	68-83 × 78-92		9, 46 - 118 × 45 - 102
Seminal receptacle size (length × width)	608 - 710	71 - 540 × 28 - 200	220 - 229 × 37-46	746-919 × 75-93	48-169 × 23-70	175-552 × 43-148	99-369 × 59-108	275-365 × 37-62	533-597 × 60-68	183 - 210 × 42 - 51	190 - 246
Egg size (length × width)	32-36 × 40-45(38- 55X38-56)	56-60 × 63-66	59-65 × 78-86	62-94 × 40-63	57-89 × 42-68	70-81 × 67-77	76-85 × 72-83	46-56 × 57-69	65-80 × 49-62	28 × 31	57 - 76 × 44 - 58
Oncosphere size (length × width)	20-33 × 22- 37(18-32 × 19-33)	23-25 × 28-31	-		23-49	38-45 × 38-40	38-50 × 34-43	22-26 × 26-35		23 × 20	40 - 57 × 31 - 53
Embryonic Hook size	11-16	14-15	16-20	14-20	17-20	13-16	16-20	17-18	16-21	12 - 18	15.7 - 20.5



Table 3

Genetic distance for specimens of *Hymenolepis folkerti* and *Hymenolepis ackerti* based on the mt locus ND1. Fractions below the diagonal represent the genetic distance based on the Jukes Cantor model. The numbers above the diagonal represent the absolute number of point mutations among the compared sequences.

Taxon, GenBank, host, collection number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
1 <i>H. folkerti</i> KX782315 ex <i>Peromyscus leucopus</i>	-	12	12	15	4	94	94	94	94	94	94	94	94	94	94	94
2 <i>H. folkerti</i> KX792199 ex <i>Tamias amoenus</i> MSB: Para: 23512	0.019	-	8	17	12	100	100	100	100	100	100	100	100	100	100	100
3 <i>H. folkerti</i> HM149295 ex <i>Thomomys</i> sp.	0.019	0.012	-	17	12	104	104	104	104	104	104	104	104	104	104	104
4 <i>H. folkerti</i> KX792195 ex <i>P. leucopus</i> MSB: Para: 23503	0.024	0.027	0.027	-	15	94	94	94	94	94	94	94	94	94	94	94
5 <i>H. folkerti</i> OP649624 ex <i>P. leucopus</i> MSB: PARA: 35312	0.006	0.019	0.019	0.024	-	97	97	97	97	97	97	97	97	97	97	97
6 <i>H. ackerti</i> OP649627 ex <i>S. hispidus</i> HWML118078	0.161	0.173	0.181	0.161	0.167	-	0	0	1	0	0	1	0	0	1	0
7 <i>H. ackerti</i> OP649634 ex <i>N. floridana</i> HWML118077	0.161	0.173	0.181	0.161	0.167	0	-	0	1	0	0	1	0	0	1	0
8 <i>H. ackerti</i> OP649626 ex <i>S. hispidus</i> HWML118076	0.161	0.173	0.181	0.161	0.167	0	0	-	1	0	0	1	0	0	1	0
9 <i>H. ackerti</i> OP649629 ex <i>S. hispidus</i> MSB: PARA: 35314	0.161	0.173	0.181	0.161	0.167	0.002	0.002	0.002	-	1	1	0	1	1	2	1
10 <i>H. ackerti</i> OP649630 ex <i>S. hispidus</i> MSB: PARA: 35319	0.161	0.173	0.181	0.161	0.167	0	0	0	0.002	-	0	1	0	0	1	0
11 <i>H. ackerti</i> NK269986 ex <i>S. hispidus</i> MSB: Mamm: 330605	0.161	0.173	0.181	0.161	0.167	0	0	0	0.002	0	-	1	0	0	1	0
12 <i>H. ackerti</i> OP649632 ex <i>S. hispidus</i> MSB: PARA: 35317	0.161	0.173	0.181	0.161	0.167	0.002	0.002	0.002	0	0.002	0.002	-	1	1	2	1
13 <i>H. ackerti</i> OP649631 ex <i>S. hispidus</i> MSB: PARA: 35318	0.161	0.173	0.181	0.161	0.167	0	0	0	0.002	0	0	0.0016	-	0	1	0
14 <i>H. ackerti</i> OP649628 ex <i>S. hispidus</i> MSB: PARA: 35313	0.161	0.173	0.181	0.161	0.167	0	0	0	0.002	0	0	0.0016	0	-	1	0
15 <i>H. ackerti</i> OP649633 ex <i>S. hispidus</i> MSB: PARA: 35316	0.161	0.173	0.181	0.161	0.167	0.002	0.002	0.002	0.003	0.002	0.002	0.0031	0.002	0.002	-	1
16 <i>H. ackerti</i> OP649625 ex <i>S. hispidus</i> MSB: PARA: 35315	0.161	0.173	0.181	0.161	0.167	0	0	0	0.002	0	0	0.0016	0	0	0.002	-

infecting hispid cotton rats is similar to the strobilar size of *H. diminuta* (about 180 mm). However, this is quite variable and it appears that it depends on the number of individuals infecting a single host. In hosts infected by only 2 parasites, worms measuring up to 107 mm (MSB: Mamm: 330681), yet for high intensity infections (~30 individuals) worms seldom exceeded 51 mm (MSB: Mamm: 305616).

Relative to the linear testicular arrangement of *H. ackerti*, it must be noted that in most of the specimens the internal testis is slightly more posterior in position compared with both poral and antiporal testes. This is a very similar arrangement as the one seen in *H. weldensis* and *H. robertrauschi*. Interestingly, the testicular arrangement in the latter is described as triangular, whereas the arrangement in the former is described as linear. Perhaps the only three species featuring an authentic triangular arrangement are *H. folkertsi*, *H. pitymi* and *H. tualatinensis*, which coincidentally also feature a rostral projection of the rostellar capsule. Yet, even in this group, there is considerable variation in the testicular arrangement, as shown by the clearly triangular arrangement of the type specimens of *H. folkertsi* (Makarikov et al., 2015), which contrast with the almost linear arrangement seen in conspecific specimens identified here (specimen collected from the prairie vole MSB: Mamm: 331573) and documented subsequently (Hoberg et al., 2016).

The phylogenetic position of parasite specimens hosted by hispid cotton rats from the Konza prairie suggests that these specimens are not conspecific with *H. diminuta* nor *H. folkertsi*, other species that occur in North America and are expected to infect a wide variety of rodents. Further, their morphological traits appear to be unique among all species known from North America, suggesting that an endemic species of *Hymenolepis* infects voles, hispid cotton rats and other rodents through the Great Plains region. Parasites were recovered from rodent hosts sampled from multiple habitat types including open grassland, shrubby habitats, and woodland. However, shrubby and woody habitats yielded the vast majority of specimens, with potential implications for increased intensity of infection through ongoing woody encroachment of the region (Briggs et al., 2005). Recent reevaluation of specimens and new helminthological surveys have resulted in the description of 4 species of hymenolepidids in the continental United States in recent years (Gardner et al., 2014; Makarikov et al., 2020, 2015). We posit that increasing sampling effort of tapeworms from rodents may yield a modest increase in the number of species described. Nevertheless, the completion of this effort will ensure a better functional understanding of the relationship between parasites, their arthropod intermediate hosts, and mammals; in turn,

setting a foundation for establishing the environmental drivers of evolution for these parasites, and variable co-evolution among multiple taxonomic groups.

Our molecular analyses also revealed a dearth of genetic information for *Hymenolepis* tapeworms in universal data repositories. From all the species endemic to North America, there are only 2 with entries in the universal repositories of the NCBI (i.e., GenBank) for the homologous gene. This lack of representation may have contributed to the evident polytomy displayed in the phylogeny based on the mitochondrial locus. To rigorously document the existing diversity and functional significance of parasites across North America, it is critical that existing personal collections, new specimen accessions, and all associated host and molecular data are deposited as voucher representation within public research archives to enable future interpretation of their distributions, host associations, and evolutionary dynamics (Colella et al., 2021; Dunnun et al., 2017; Galbreath et al., 2019).

This practice is particularly important for *Hymenolepis*, a group that includes species of zoonotic importance, such as the cosmopolitan *H. diminuta* (Panti-May et al., 2020). Evidence gathered elsewhere suggests that there may be several species in the genus that have a zoonotic potential (Nkouawa et al., 2016), which may have been identified as *H. diminuta* based on the limited variability of characters available in the specimens used for the parasite diagnosis. In most cases these specimens include eggs and meristic characters as evidence for species level identification, yet there is overlap in the range of these measurements, such as the overall size of the embryophore and hooks. As demonstrated by Nkouawa et al. (2016), molecular screening allows a more precise identification of the specimens, which is not possible with a limited set of overlapping characters. Their results may suggest that some infections attributed to *H. diminuta* in humans may be induced by other species. As indicated by Panti-May et al. (2020) these identifications are based on the morphology of the eggs and not on the morphology of the adults. As seen in Table 3, the eggs feature very few meristic characters that overlap for several species.

This would be an unfortunate example of the “fallacy of the expected identification” (Hoberg & Soudachanh, 2021), one with important medical implications. In this phenomenon, scientists assign a name of a species that has been recorded in the host species elsewhere without the proper verification of the identity based on the analysis of the voucher specimen or DNA. Typically, this species is presumed common and widespread. We plea to the scientific community to properly voucher the material, which may allow other members of the community to evaluate it and offer a refined identification. An example of

this practice includes the deposition of specimens initially identified as *H. diminuta* and collected from cotton rats (Faulkner & Lochmiller, 2000), examination of these specimens deposited in the USNM1382519 revealed that they belong to a different species. Another example includes the proper vouchering of specimens and the generation of barcodes accessible to the community (Sasaki et al., 2021). We propose that whenever possible, scientist excise tissues from specimens that will be eventually stained for identification and deposition in scientific collections.

Tapeworms are likely valuable indicators of biodiversity change across both geography and time, considering they require at least one arthropod intermediate host (Gardner & Campbell, 1992). Fluctuations in parasite abundance, distribution, and host association should prove important to our understanding of general biotic responses to anthropogenic environmental change.

## Acknowledgements

Kansas State Division of Biology and the Department of Biology supported AGH, NSF-DUE 1564969 supported the writing of this description. Sara Brant (Museum of Southwestern Biology, University of New Mexico), Anna Phillips and Lisa Comer (Smithsonian Institution), Luis García Prieto (Colección Nacional de Helminthos, UNAM), and Gabor Racz (Harold W. Manter Laboratory, University of Nebraska) made arrangements for the examination and deposition of the helminthological material.

## References

- Briggs, J. M., Knapp, A. K., Blair, J. M., Heisler, J. L., Hoch, G. A., Lett, M. S. et al. (2005). An ecosystem in transition. Causes and consequences of the conversion of mesic grassland to shrubland. *Bioscience*, 55, 243–254.
- Bruckerhoff, L., Connell, K., Guinnip, J., Adhikari, E., Godar, A., Gido, K. et al. (2021). Harmony on the prairie? Grassland plant and animal community responses to variation in climate across land-use gradients. *Ecology*, 102, e02986. <https://doi.org/10.1002/ecy.2986>
- Coggins, J. R., & McDaniel, J. S. (1975). Helminth population dynamics in the cotton rat, *Sigmodon hispidus*. *Proceedings of the Oklahoma Academy of Science*, 55, 112–118.
- Colella, J. P., Bates, J., Burneo, S. F., Camacho, M. A., Bonilla, C. C., Constable et al. (2021). Leveraging natural history biorepositories as a global, decentralized, pathogen surveillance network. *Plos Pathogens*, 17, e1009583. <https://doi.org/10.1371/journal.ppat.1009583>
- Dunnum, J. L., Yanagihara, R., Johnson, K. M., Armien, B., Batsaikhan, N., Morgan, L. et al. (2017). Biospecimen Repositories and Integrated Databases as Critical Infrastructure for Pathogen Discovery and Pathobiology Research. *Plos Neglected Tropical Diseases*, 11, e0005133. <https://doi.org/10.1371/journal.pntd.0005133>
- Faulkner, B. C., & Lochmiller, R. L. (2000). Ecotoxicity revealed in parasite communities of *Sigmodon hispidus* in terrestrial environments contaminated with petrochemicals. *Environmental Pollution*, 110, 135–145. [https://doi.org/10.1016/S0269-7491\(99\)00276-6](https://doi.org/10.1016/S0269-7491(99)00276-6)
- Galbreath, K. E., Hoberg, E. P., Cook J. A., Armien, B., Bell, K. C., Campbell, M. L. et al. (2019). Building an integrated infrastructure for exploring biodiversity: field collections and archives of mammals and parasites. *Journal of Mammalogy*, 100, 382–393. <https://doi.org/10.1093/jmammal/gyz048>
- Gardner, S. L. (1985). Helminth parasites of *Thomomys bulbivorus* (Richardson) (Rodentia: Geomyidae), with the description of a new species of *Hymenolepis* (Cestoda). *Canadian Journal of Zoology*, 63, 1463–1469.
- Gardner, S. L., & Campbell, M. L. (1992). Parasites as probes for biodiversity. *Journal of Parasitology*, 78, 596–600.
- Gardner, S. L., Dursahinhan, A. T., Campbell, M. L., & Racz, S. E. (2020). A new genus and two new species of unarmed hymenolepidid cestodes (Cestoda: Hymenolepididae) from geomyid rodents in Mexico and Costa Rica. *Zootaxa*, 4766, 358–376. <https://doi.org/10.11646/zootaxa.4766.2.5>
- Gardner, S. L., Luedders, B. A., & Duszynski, D. W. (2014). *Hymenolepis robertrauschi* n. sp. from grasshopper mice *Onychomys* spp. in New Mexico and Nebraska, U.S.A. *Occasional Papers of the Texas Tech University Museum*, 322, 1–10.
- Harkema, R. (1946). The Metazoa parasitic in cotton rats of Wake County. *Journal of the Elisha Mitchell Scientific Society*, 62, 142–143.
- Harkema, R., & Kartman, L. (1948). Observations on the helminths and ectoparasites of the cotton rat, *Sigmodon hispidus* Say and Ord, in Georgia and North Carolina. *Journal of the Elisha Mitchell Scientific Society*, 64, 183–191.
- Haukisalmi, V., Hardman, L. M., Foronda, P., Feliu, C., & Henttonen, H. (2010). Systematic relationships of *Mosgovoyia* Spasskii, 1951 (Cestoda: Anoplocephalidae) and related genera inferred from mitochondrial and nuclear sequence data. *Systematic Parasitology*, 77, 71–79.
- Hoberg, E. P., Makarikov, A. A., Tkach, V. V., Meagher, S., Nims, T. N., Eckerlin, R. P. et al. (2016). Insights on the host associations and geographic distribution of *Hymenolepis folkertsi* (Cestoda: Hymenolepididae) among rodents across temperate latitudes of North America. *Parasitology Research*, 115, 4627–4638.
- Hoberg, E. P., & Soudachanh, K. M. (2021). Diversity of Tetrabothriidae (Eucestoda) among Holarctic Alcidae (Charadriiformes): Resolution of the *Tetrabothrius jagerskioeldi* cryptic species complex —cestodes of Alcinae— provides insights on the dynamic nature of tapeworm and marine bird faunas under the Stockholm Paradigm. *MANTER. Journal of Parasite Biodiversity*, 16, 1–76. <https://doi.org/10.32873/unl.dc.manter16>
- Hope, A. G. (2019). *Small mammal host-parasite sampling data for 16 linear trapping transects located in 8 LTER*

- burn treatment watersheds at Konza Prairie version 3. Environmental Data Initiative. <https://doi.org/10.6073/pasta/69109c56fcf21a30a8d37369cb47f8de>
- Makarikov, A. A., Galbreath, K. E., Eckerlin, R. P., & Hoberg, E. (2020). Discovery of *Arostrilepis* tapeworms (Cyclophyllidae: Hymenolepididae) and new insights for parasite species diversity from Eastern North America. *Parasitology Research*, *119*, 567–585. <https://doi.org/https://doi.org/10.1007/s00436-019-06584-4>
- Makarikov, A. A., Nims, T. N., Galbreath, K. E., & Hoberg, E. P. (2015). *Hymenolepis folkertsi* n. sp (Eucestoda: Hymenolepididae) in the oldfield mouse *Peromyscus polionotus* (Wagner) (Rodentia: Cricetidae: Neotominae) from the southeastern Nearctic with comments on tapeworm faunal diversity among deer mice. *Parasitology Research*, *114*, 2107–2117. <https://doi.org/10.1007/s00436-015-4399-x>
- Miller, M. A., Pfeiffer, W., & Schwartz, T. (2010). Creating the CIPRES Science Gateway for inference of large phylogenetic trees. New Orleans: Gateway Computing Environments Workshop (GCE). <https://doi.org/10.1109/GCE.2010.5676129>
- Mollhagan, T. (1978). Habitat influence on helminth parasitism of the cotton rat in western Texas with remarks on some of the parasites [Article]. *Southwestern Naturalist*, *23*, 401–408. <https://doi.org/10.2307/3670248>
- Morgan, J. A. T., & Blair, D. (1998). Relative merits of nuclear ribosomal internal transcribed spacers and mitochondrial CO1 and ND1 genes for distinguishing among *Echinostoma* species (Trematoda). *Parasitology*, *116*, 289–297.
- Nkouawa, A., Haukisalmi, V., Li, T. Y., Nakao, M., Lavikainen, A., Chen, X. W. et al. (2016). Cryptic diversity in hymenolepidid tapeworms infecting humans. *Parasitology International*, *65*, 83–86. <https://doi.org/10.1016/j.parint.2015.10.009>
- Panti-May, J. A., Rodríguez-Vivas, R. I., García-Prieto, L., Servian, A., & Costa, F. (2020). Worldwide overview of human infections with *Hymenolepis diminuta*. *Parasitology Research*, *119*, 1997–2004. <https://doi.org/10.1007/s00436-020-06663-x>
- Pfaffenberger, G. S., Kemether, K., & Debruin, D. (1985). Helminths of sympatric populations of kangaroo rats (*Dipodomys ordii*) and grasshopper mice (*Onychomys leucogaster*) from the high plains of eastern New Mexico. *Journal of Parasitology*, *71*, 592–595. <https://doi.org/10.2307/3281429>
- Rambaut, A. (2012). FigTree v1. 4. *Molecular evolution, phylogenetics and epidemiology*. Edinburgh: University of Edinburgh, Institute of Evolutionary Biology.
- Ratajczak, Z., Briggs, J. M., Goodin, D. G., Luo, L., Mohler, R. L., Nippert, J. B. et al. (2016). Assessing the potential for transitions from Tallgrass Prairie to Woodlands: are we operating beyond critical fire thresholds? *Rangeland Ecology & Management*, *69*, 280–287. <https://doi.org/10.1016/j.rama.2016.03.004>
- Riley, W. A., & Shannon, W. R. (1922). The rat tapeworm, *Hymenolepis diminuta*, in man. *The Journal of Parasitology*, *8*, 109–117. <https://doi.org/10.2307/3270923>
- Sasaki, M., Anders, J. L., & Nakao, M. (2021). Cestode fauna of murid and cricetid rodents in Hokkaido, Japan, with assignment of DNA barcodes. *Species Diversity*, *26*, 255–272. <https://doi.org/10.12782/specdiv.26.255>
- Seidenberg, A. J., Kelly, P. C., Lubin, E. R., & Buffington, J. D. (1974). Helminths of cotton rat in southern Virginia, with comments on sex-ratios of parasitic nematode populations [Article]. *American Midland Naturalist*, *92*, 320–326. <https://doi.org/10.2307/2424297>
- Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W. et al. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*, *7*, 539.
- Smith, C. S., Jones, F. E., & Eyles, D. E. (1953). Three additional Tennessee cases of *Hymenolepis diminuta* infection in man. *Journal of the Tennessee Academy of Science*, *28*, 196–197.